

Complementarity between Gauge-Boson Compositeness and Asymptotic Freedom

Keiichi Akama

Department of Physics, Saitama Medical College, Kawakado, Moroyama, Saitama, 350-04, Japan

Takashi Hattori

Department of Physics, Kanagawa Dental College, Inaoka-cho, Yokosuka, Kanagawa, 238, Japan

We derive and solve the compositeness condition for the $SU(N_c)$ gauge boson at the next-to-leading order in $1/N_f$ (N_f is the number of flavors) and the leading order in $\ln \Lambda^2$ (Λ is the compositeness scale) to obtain an expression for the gauge coupling constant in terms of the compositeness scale. It turns out that the argument of gauge-boson compositeness (with a large Λ) is successful only when $N_f/N_c > 11/2$, in which the asymptotic freedom fails.

PACS number(s): 12.60.Rc, 11.10.Jj, 11.15.Pg, 11.15.-q

Recently, the possible compositeness of gauge bosons [1–6] has attracted renewed attention from both theoretical [7–10] and phenomenological [11,12] points of view. The experimental results at present do not exclude [13] and may even suggest [11] the interesting possibility that quarks, leptons and gauge bosons are composite [12]. Based on general theoretical analyses [2,7], this idea has been widely applied in various branches of physics. In quark-lepton physics, various models have been considered in terms of composite gauge bosons [3,8]. In hadron physics, the vector mesons can be interpreted as gauge bosons with hidden local symmetry [4,9]. The gauge fields induced in connection with the geometric phase (Berry phase) in molecular and other systems are also expected to become dynamical through the quantum effects of matter, and could be considered as composite gauge bosons [5,6,10].

A gauge boson interacting with matter can be interpreted as composite under the compositeness condition $Z_3 = 0$ [14], where Z_3 is the wave-function renormalization constant of the gauge boson. Under this condition, the gauge field becomes an auxiliary field without independent degrees of freedom [15]. The quantum fluctuations, however, give rise to a kinetic term of the gauge field, so that a dynamical gauge boson is induced as a composite of the matter fields. The compositeness conditions which have been investigated so far hold only in the large N limit, where N is the number of the matter fields coupling with the gauge boson. However, in cases of practical interest, N is rather small. Thus it is important to investigate the higher order effects in $1/N$. In our previous papers, we derived the next-to-leading order (NLO) contributions to the compositeness conditions in the Nambu-Jona-Lasinio model [16] and the abelian gauge theory [17] at the leading order in $\ln(\Lambda^2/m^2)$ (Λ is the compositeness scale, and m is the constituent mass scale [18]). In this paper, we perform a similar investigation for the non-abelian gauge theory, and find that the argument of gauge-boson compositeness (with a large Λ) is successful only when the number of flavors, N_f , is so large that the gauge theory is not asymptotically free [19]. On the other hand, many people have argued that the asymptotically non-free theory may encounter the problem of excessively large coupling constants at some ultraviolet energy scale, suggesting the necessity for some new physics such as compositeness [20]. Thus we observe a complementarity between gauge-boson compositeness (with a large Λ), if any, and asymptotic freedom in the gauge theories.

We consider the $SU(N_c)$ gauge theory for the gauge boson G_μ^a ($a = 1, \dots, N_c^2 - 1$) with N_f fermionic matters ψ_j ($j = 1, \dots, N_f$):

$$\begin{aligned} \mathcal{L}^{(\Lambda)} = & -\frac{1}{4} (G_{\mu\nu}^a)^2 + \sum_j \bar{\psi}_j \left(i \not{\partial} - m + \frac{1}{2} g \lambda^a \not{G}^a \right) \psi_j \\ & - \frac{1}{2\alpha} (\partial^\mu G_\mu^a)^2 + \partial^\mu \eta^{a\dagger} (\partial_\mu \eta^a - g f^{abc} \eta^b G_\mu^c) \end{aligned} \quad (1)$$

where $G_{\mu\nu}^a = \partial_\mu G_\nu^a - \partial_\nu G_\mu^a + g f^{abc} G_\mu^b G_\nu^c$, g is the coupling constant, m is the mass of ψ_j , α is the gauge fixing parameter, and η^a ($a = 1, \dots, N_c^2 - 1$) is the Fadeev-Popov ghost. The superscript (Λ) of $\mathcal{L}$ indicates that the bare theory has a large but finite ultraviolet cutoff Λ . In order to absorb the ultraviolet divergences arising from quantum fluctuations, we renormalize the fields and coupling constant as [21]

$$\begin{aligned} \mathcal{L}^{(\Lambda)} = & -\frac{1}{4} Z_3 (G_{r\mu\nu}^a)^2 + \sum_j \bar{\psi}_{rj} \left(i Z_2 \not{\partial} - Z_m m_r + \frac{1}{2} Z_1 g_r \lambda^a \not{G}_r^a \right) \psi_{rj} \\ & - \frac{Z_3}{2Z_\alpha \alpha_r} (\partial^\mu G_{r\mu}^a)^2 + Z_\eta \partial^\mu \eta_r^{a\dagger} \left(\partial_\mu \eta_r^a - \frac{Z_1}{Z_2} g_r f^{abc} \eta_r^b G_{r\mu}^c \right) \end{aligned} \quad (2)$$

where $G_{r\mu\nu}^i = \partial_\mu G_{r\nu}^a - \partial_\nu G_{r\mu}^a + (Z_1/Z_2)g_r f^{abc}G_{r\mu}^b G_{r\nu}^c$, the quantities with the index “r” are the renormalized ones, and $Z_1, Z_2, Z_3, Z_\eta, Z_m$, and Z_α are the renormalization constants.

Now let us impose the condition

$$Z_3 = 0. \quad (3)$$

Under the condition (3), Lagrangian (2) involves no derivative of $G_{r\mu}^i$ so that the Euler equation with respect to $G_{r\mu}^i$ becomes a constraint, and $G_{r\mu}^i$ is an auxiliary field without independent dynamical degrees of freedom. To get the physical spectrum, however, we need to follow the usual renormalization procedure with (2), where the physical gauge boson states do exist even under the condition (3). This means that the gauge boson is induced as a composite of matter fields, and hence (3) is called “compositeness condition” [14].

At the lowest order in g_r^2 , Z_3 is determined so as to cancel the divergence in the one-loop diagrams in Fig. A–C. As is well known, this is given by [21]

$$Z_3 = 1 - \left[\frac{2}{3}N_f - \left(\frac{13}{6} - \frac{\alpha_r}{2} \right) N_c \right] g_r^2 I, \quad (4)$$

where I is the divergent integral, given in the dimensional regularization by $I = 1/16\pi^2\epsilon = 1/8\pi^2(4-d)$ with the number of spacetime dimensions d . A simple-minded application of the compositeness condition $Z_3 = 0$ gives

$$g_r^2 = 1 / \left[\frac{2}{3}N_f - \left(\frac{13}{6} - \frac{\alpha_r}{2} \right) N_c \right] I. \quad (5)$$

To avoid the absurdity of a vanishing coupling constant, we take the regularization as an approximation to some physical cutoff, and fix ϵ at the non-vanishing value $\epsilon = 1/\ln(\Lambda^2/m^2)$. If we take the relation (5) as the leading one, infinitely many diagrams of any higher order in g_r have the same order of magnitude, and therefore the usual perturbation expansion in g_r fails, as in the cases investigated in our previous works [16,17]. Then we need to rely on a $1/N_f$ or $1/N_c$ expansion. If N_c is large, the solution for g_r^2 in (5) becomes negative, and worse, infinitely many higher loops with internal gauge-boson lines would also belong to the leading order in $1/N_c$. Thus, the expansion in $1/N_c$ is not appropriate. On the other hand, $1/N_f$ can successfully classify diagrams, as we shall see below. In this case, the leading order contribution comes from the one-fermion-loop diagram (Fig. A), and the one-boson-loop diagrams (Fig. B,C) belong to the next-to-leading order. This harmonizes well with the physical picture that the gauge boson is a composite of the fermions.

Now we turn to the NLO contributions in $1/N_f$. In addition to the one-boson-loop diagrams B and C in Fig. , the multi-loop diagrams D–H in Fig. belong to this order. In Fig. , the line of small circles stands for the gauge boson propagator with an arbitrary number of one-fermion-loops inserted. The renormalization constant Z_3 should be chosen so as to cancel all the divergence in these diagrams. Though the n -loop diagram diverges like I^n , it is suppressed by the factor $g^{2n} \propto 1/I^n$. Therefore, the leading divergent part of each diagram is $O(I^n)O(I^{-n}) \sim O(1)$, apart from the powers of $1/N_f$. We denote this leading contribution of $O(1)$ from diagram X (=D–H in Fig.) to Z_3 by $Z(X, l)$, where l is the number of the fermion loops per diagram. Different diagrams with the same X and l give $Z(X, l)$ exactly the same contributions. We call the number of such different diagrams “multiplicity” of X with l . The overlapping divergence in G and H is separated into two parts: “f” in respect to the fermion loop divergence at the three boson vertex part, and “m” in respect to the boson-fermion (mixed) loop divergence at the boson-fermion-fermion vertex part. We denote these contributions as $Z(\text{Gf}, l)$, $Z(\text{Gm}, l)$, $Z(\text{Hf}, l)$, and $Z(\text{Hm}, l)$.

The following properties are useful in the calculation. (a) Due to the gauge symmetry, the leading divergences cancel each other in two diagrams: that with a fermion-loop inserted at a three-gauge-boson vertex in some diagram and that with a fermion-loop inserted into a gauge-boson propagator adjacent to the vertex. (b) Because the one-fermion-loop (denoted by $\Pi_0^{\mu\nu}$) inserted into a gauge-boson propagator (with momentum q) is divergenceless (i.e. $q_\mu \Pi_0^{\mu\nu} = 0$), it diminishes the α_r -dependent part of the diagram.

The divergent contribution $Z(X, l)$ is separated into the gauge independent part $Z_0(X, l)$ and the part $Z_\alpha(X, l)$ linear in α_r , while the terms higher in α_r are convergent. We first consider the α_r -independent parts. Because the lowest-order fermion self-energy part and boson-fermion-fermion vertex part converge in the Landau gauge, $Z_0(\text{D}, l)$ and $Z_0(\text{E}, l)$ have no leading divergence. Because property (a) implies $Z_0(\text{F}, l) = -Z_0(\text{Gf}, l) = Z_0(\text{Hf}, l)$ for relevant l , and their multiplicities are $l+1$, $2l$, and $l-1$, respectively, they cancel out when $l \geq 1$. For $l=0$, diagrams G and H are absent, while diagram F is nothing but diagram B (including the α_r -dependent part), and, together with diagram C, contributes the second term in the square brackets in (4). Because property (a) implies $Z_0(\text{Gm}, l) = -Z_0(\text{Hm}, l)$ ($l \geq 2$) and their multiplicities are $2l$ and $2(l-1)$, respectively, diagrams Gm and Hm contribute the following terms to Z_3 :

$$-\sum_{l=1}^{\infty} \frac{1}{l(l+1)} \left(-\frac{2}{3}\right)^{l-1} N_c N_f^l (g_r^2 I)^{l+1}. \quad (6)$$

Then we consider the α_r -dependent parts. Due to property (b), diagrams D and E have contributions only when $l = 1$, and F, G, and H have contributions only when no fermion-loop is inserted on one of the two gauge-boson lines. The multiplicities of F, Gf, and Hf, are 1, 2, and 1, respectively, and those of Gm and Hm are 2 and 2, respectively, for any relevant l . Because property (a) implies $Z_\alpha(F, l) = -Z_\alpha(Gf, l) = Z_\alpha(Hf, l)$ and $Z_\alpha(Gm, l) = -Z_\alpha(Hm, l)$ for relevant l , they cancel out for $l \geq 2$. For $l = 1$, diagram H is absent, and the contributions from diagrams D, E, F, Gf, and Gm with $l = 1$ to Z_3 are

$$-\frac{1}{3} \alpha_r N_c N_f (g_r^2 I)^2 \quad (7)$$

times $-1 + 1/N_c^2$, $-1/N_c^2$, $-1/2$, 1, and $3/2$, respectively, and hence (7) times 1 in total. The contribution from $l = 0$ has already been taken into account in (4). Next, we renormalize the subdiagram divergences by subtracting the divergent counter parts of (i) each fermion loop inserted into the gauge boson lines in D–H, (ii) the fermion self-energy part in D, (iii) the fermion-boson vertex part in E, (iv) the three-boson vertex part in Gf and Hf, and (v) the boson-fermion-fermion vertex part in Gm and Hm. The contributions from the counter parts cancel out for even l , and amount to minus twice the original terms for odd l . Thus in total they contribute the following terms to Z_3 :

$$\sum_{l=1}^{\infty} \frac{1}{l(l+1)} \left(\frac{2}{3}\right)^{l-1} N_c N_f^l (g_r^2 I)^{l+1} + \frac{1}{3} \alpha_r N_c N_f (g_r^2 I)^2. \quad (8)$$

Collecting all the contributions in (4) and (8) together, we finally obtain the compact expression

$$Z_3 = 1 - \frac{2}{3} N_f g_r^2 I + \frac{11}{3} N_c g_r^2 I - \frac{\alpha_r}{2} N_c g_r^2 I \left(1 - \frac{2}{3} N_f g_r^2 I\right) + \frac{3}{2} N_c \left(\frac{3}{2N_f} - g_r^2 I\right) \ln\left(1 - \frac{2}{3} N_f g_r^2 I\right) + O\left(\frac{1}{N_f^3}\right). \quad (9)$$

The compositeness condition $Z_3 = 0$ with expression (9) looks somewhat complex. We can, however, solve it for g_r^2 by iterating the leading-order solution into itself. The solution is rather simple:

$$g_r^2 = \frac{3}{2N_f I} \left[1 + \frac{11N_c}{2N_f} + O\left(\frac{1}{N_f^2}\right) \right]. \quad (10)$$

The logarithmic term in (9) is suppressed in the solution (10) by the factor which vanishes with iteration of the leading solution. It is interesting that the solution does not depend on the gauge parameter α_r , in spite of the fact that Z_3 does. The α_r -dependent term in (9) is also suppressed in the solution in the same way that the logarithmic term is. The solution of the compositeness condition should be gauge-independent because it is a relation among physically observable quantities. Note that we assumed a large but finite physical cutoff Λ , by fixing ϵ at a non-vanishing value $\epsilon = 1/\ln(\Lambda^2/m^2)$. Because the above argument relies on $1/N_f$ expansion including iteration, the absolute value of the next-to-leading contribution should not exceed that of the leading one. If we apply it to (10), we obtain

$$N_f > \frac{11N_c}{2}. \quad (11)$$

The allowed region of N_f/N_c by (11) is complementary to that for asymptotic freedom in the gauge theory [19]. When the gauge theory is asymptotically free, the next-to-leading contributions to the compositeness condition are so large that the gauge bosons cannot be composites of the above type. On the contrary, when the theory is asymptotically non-free, the next-to-leading order contributions are suppressed, and the gauge bosons can be interpreted as composite [22]. As was shown in the previous paper [17], the NLO contribution is suppressed in the abelian gauge theory, making the compositeness interpretation successful. This is in accordance with the above-mentioned complementarity, since the abelian gauge theory is not asymptotically free. In general, in the asymptotically non-free theories with the beta function derived via perturbations in the coupling constant, the running coupling constant diverges at some ultra-violet scale. Then we should introduce a momentum cutoff, since the theory gives nothing above that scale. Some new physics such as compositeness is required to fill in this blank [20].

The marginal value $N_f/N_c = 11/2$ for asymptotic freedom was derived via the renormalization group method at the leading order in g_r^2 (and exactly in N_c and N_f). All the one-loop diagrams and the renormalization constants Z_1 , Z_2 , and Z_3 were used there. On the other hand, (11) was derived via the NLO perturbation in $1/N_f$, and by using multi-loop boson-self-energy diagrams and Z_3 only. The coincidence of the marginal values $N_f/N_c = 11/2$ may not be accidental, however, because in the latter the large- Λ approximation implied small g_r , and we implicitly used the other one-loop diagrams (other than the boson self-energy) as subdiagrams, and also used Z_1 and Z_2 (besides Z_3) to renormalize the divergences due to the subdiagrams, while in the former (the renormalization group method) one-loop contributions at some scale could be interpreted as a sum of multi-loops at the other scale. It would be interesting to confirm this complementarity at a higher order, or in other models such as those with scalar matter, or to argue from a more general point of view covering all orders. We expect that these methods and results concerning NLO compositeness and the concept of complementarity presented here will be useful in pursuing the composite dynamics of nuclei and hadrons, and the possible compositeness of quarks, leptons, gauge bosons and Higgs scalars, as well as useful in induced gauge theories concerning molecular, solid-state, and other systems.

We would like to thank Professor H. Terazawa for useful discussions.

-
- [1] W. Heisenberg, *Rev. Mod. Phys.* **29**, 269 (1957); J. D. Bjorken, *Ann. Phys.* **24**, 174 (1963).
 - [2] I. Bialynicki-Birula, *Phys. Rev.* **130**, 465 (1963); D. Lurié and A. J. Macfarlane, *Phys. Rev.* **136** (1964) B816; T. Eguchi, *Phys. Rev.* **D14** (1976) 2755; **D17** (1978) 611; K. Shizuya, *Phys. Rev.* **D21** (1980) 2327.
 - [3] K. Akama and H. Terazawa, *INS-Rep-257* (1976); H. Terazawa, Y. Chikashige and K. Akama, *Phys. Rev.* **D15**, 480 (1977); T. Saito and K. Shigemoto, *Prog. Theor. Phys.* **57** (1977) 242; O. W. Greenberg and J. Sucher, *Phys. Lett.* **99B**, 339 (1981); L. F. Abbott and E. Farhi, *Phys. Lett.* **101B**, 69 (1981).
 - [4] See for example, J. J. Sakurai, *Currents and Mesons*, (Univ. Chicago Press, Chicago, 1969); M. Bando, T. Kugo, S. Uehara, K. Yamawaki and T. Yanagida, *Phys. Rev. Lett.* **54** 1215 (1985).
 - [5] C. A. Mead and D. G. Truhlar, *J. Chem. Phys.* **70(05)**, 2284 (1979); M. V. Berry, *Proc. R. Soc. London Ser.* **A392**, 45 (1984).
 - [6] K. Akama, in *Gauge Theory and Gravitation*, Proceedings of the International Symposium, Nara, Japan, 1982 ed. K. Kikkawa, N. Nakanishi and H. Nariai (Springer-Verlag) p. 267; M. D. Maia, *J. Math. Phys.* **25** (1984) 2090; K. Akama, *Prog. Theor. Phys.* **78**, 184 (1987); **79**, 1299 (1988).
 - [7] A. Hasenfratz and P. Hasenfratz, *Phys. Lett.* **B297**, 166 (1992); M. A. Luty, *Phys. Rev.* **D48**, 1295 (1993); K. Akama and T. Hattori, *Phys. Lett.* **B392**, 383 (1997); M. Bando, Y. Taniguchi and S. Tanimura, *Prog. Theor. Phys.* **97**, 665 (1997); M. Bando, J. Sato, and K. Yoshioka, *ICRR-Report-384-97-7*, hep-ph/9703321 (1997); M. Bando, hep-ph/9705237 (1997).
 - [8] K. Akama and T. Hattori, *Int. J. Mod. Phys.* **A9**, 3503 (1994); B. S. Balakrishna and K. T. Mahanthappa, *Phys. Rev.* **D49** 2653 (1994); *Phys. Rev.* **D52** 2379 (1995); A. Galli, *Phys. Rev.* **D51**, 3876 (1995); *Nucl. Phys.* **B435**, 339 (1995); C. P. Burgess, J.-P. Derendinger, F. Quevedo, and M. Quiros, *Phys. Lett.* **B348** 428 (1995); V. V. Kabachenko and Y. F. Pirogov, *IHEP 96-97*, hep-ph/9612275 (1996); M. Berkooz, P. Cho, P. Kraus, and M. J. Strassler, *HUTP-97/A014*, hep-th/9705003 (1997); M. Yasuè, *TOKAI-HEP/TH-9701*, hep-ph/9707312 (1997).
 - [9] M. Tanabashi, *Phys. Lett.* **B384**, 218 (1996); M. C. Birse, *Z. Phys.* **A355**, 231 (1996); M. Hashimoto, *Phys. Rev.* **D54**, 5619 (1996); C. M. Shakin and W.-D. Sun, *Phys. Rev.* **D55**, 2874 (1997); L.-H. Chan, *Phys. Rev.* **D55**, 5362 (1997); M. Harada and A. Shibata, *Phys. Rev.* **D55**, 6716 (1997).
 - [10] K. Kikkawa, *Phys. Lett.* **B297**, 89 (1992); T. Isse, *Mod. Phys. Lett.* **A8** (1993) 837, 947; T. Hatsuda and H. Kuratsuji, *UTHEP-286* (1994); K. Kikkawa and H. Tamura, *Int. J. Mod. Phys.* **A10**, 1597 (1995); S. Tanimura and I. Tsutsui, *Mod. Phys. Lett.* **A10**, 2607 (1995); E. M. Monte and M. D. Maia, *UNB-MAT-FM-M495* (1995).
 - [11] CDF Collaboration, F. Abe *et al.*, *Phys. Rev. Lett.* **77**, 438 (1996); *ibid.* **77**, 5336 (1996); *Phys. Rev. D* **55**, R5263 (1997); H1 Collaboration, C. Adloff *et al.*, *Z. Phys. C* **74**, 191 (1997); ZEUS Collaboration, J. Breitweg *et al.*, *Z. Phys. C* **74**, 207 (1997).
 - [12] M. Bander, *Phys. Rev. Lett.* **77**, 601 (1996); K. Akama and H. Terazawa, *Phys. Rev. D* **55**, R2521 (1997); A. E. Nelson, *Phys. Rev. Lett.* **78** 4159 (1997); K. Akama K. Katsuura, and H. Terazawa, *Phys. Rev. D* **56**, R2490 (1997); S. L. Adler, *IASSNS-HEP 97-12*, hep-ph/9702378; M. C. Gonzalez-Garcia and S.F. Novaes, *IFT-P-024-97*, hep-ph/9703346; N. Di Bartolomeo and M. Fabbrichesi, *Phys. Lett.* **B406**, 237 (1997).
 - [13] Particle Data Group, R.M. Barnett *et al.*, *Phys. Rev. D* **54**, 1(1996).
 - [14] B. Jouvet, *Nuovo Cim.* **5**, 1133 (1956); M. T. Vaughn, R. Aaron and R. D. Amado, *Phys. Rev.* **124**, 1258 (1961); A. Salam, *Nuovo Cim.* **25**, 224 (1962); S. Weinberg, *Phys. Rev.* **130**, 776 (1963); Lurié and Macfarlane, *Ref. [2]*; Eguchi, *Ref. [2]*; Shizuya, *Ref. [2]*.
 - [15] D. J. Gross and A. Neveu, *Phys. Rev.* **D10**, 3235 (1974); T. Kugo, *Prog. Theor. Phys.* **55**, 2032 (1976); T. Kikkawa, *Prog. Theor. Phys.* **56**, 947 (1976).

- [16] K. Akama, Phys. Rev. Lett. **76**, 184 (1996).
- [17] Akama and Hattori, Ref. [7].
- [18] In practice, we use the dimensional regularization with correspondence $\epsilon = (4 - d)/2 = 1/\ln(\Lambda^2/m^2)$, where d is the number of spacetime dimensions.
- [19] D. J. Gross and F. Wilczek, Phys. Rev. Lett. **30**, 1343 (1973); H. D. Politzer, Phys. Rev. Lett. **30**, 1346 (1973).
- [20] W.A. Bardeen, C.T. Hill and M. Lindner, Phys. Rev. **D41**, 1647 (1990); M. Harada, Y. Kikukawa, T. Kugo and H. Nakano, Prog. Theor. Phys. **92**, 1161 (1994); Bando, Taniguchi and Tanimura, Ref. [7]; Bando, Sato and Yoshioka, Ref. [7]; Bando, Ref. [7].
- [21] See for example, T. Muta, *Foundations of Quantum Chromodynamics, An Introduction to Perturbative Method in Gauge Theories*, World Scientific, §2.5.5. (1987).
- [22] This does not eliminate the possibility that the composite gauge boson with a low Λ is compatible with asymptotic freedom.




FIG. 1

The gauge-boson self-energy parts at the leading order (A) and at the next-to-leading order (B–H) in $1/N_f$. The solid, dotted, and small-dotted lines indicate the fermion, gauge-boson, and Fadeev-Popov ghost propagators, respectively. The line of small circles stands for the gauge boson propagator with an arbitrary number of fermion-loops inserted.